

Lettuce Management Schedule

*A guide to weed, insect and
disease management in field
lettuce in Nova Scotia*



2019



GUIDE TO PEST MANAGEMENT IN FIELD LETTUCE



Nova Scotia Vegetable Crop Guide to Pest Management 2019
[LET1-19]

Updated April 22, 2019 by
Peter Burgess, Horticulturist, Perennia
Mélanie Leclerc, Research Associate, Perennia

IMPORTANT

This publication was compiled by representatives from Perennia using information from the Pest Management Regulatory Agency of Health Canada, specific pesticide labels, previous Atlantic Provinces Vegetable Pest Guides and manufacturer's information. **This information is continuously changing and therefore it can cease to be current and accurate. Pesticide labels are the best source of information and should always be consulted prior to using a product.** The label is the best source of information on: registered crop uses, rates, days to harvest, compatibility with other pesticides, toxicity and other special information on its effective and safe use

By printing this publication, Perennia does not offer any warranty or guarantee, nor do they assume any liability for any crop loss, animal loss, health, safety or environmental hazard caused by the use of a pesticide mentioned in this publication.

WARNINGS

This publication is meant to be used as a reference for possible pest control options. Where there are multiple brand names of a specific active ingredient registered in Canada, Perennia has only listed a couple for reference purposes and as such does not endorse one brand over another. If you have purchased a generic product not specifically in this guide but has your crop and pest on the label, always follow that product label.

If any information in this or any other publication conflicts with the information on the label, always use the label recommendation. If you have an old label, your pesticide supplier should be able to give you the newest label. You are legally responsible for the safe use of pesticides you purchase. This means the safe transport and storage of these materials, the label rates used on crops, and the safe disposal of containers.

Pest	Group	Active Ingredient	Pesticide Product Name	Rate	Restricted Entry Interval (REI)	Pre-harvest Interval (days)	Remarks
WEEDS:							
Preplant <i>Perennial weeds including quackgrass</i>	9	glyphosate	Roundup 356 Sn	1.25-2.5 L/ha	-	7	Apply in the fall or spring prior to planting. Annual weed control programs will be necessary to control weeds germinating after planting. For quackgrass control, apply to actively growing quackgrass when at least 4 new leaves are present. The low rate (2.5 L/ha) will provide a minimum of one season control, while higher rates (4.75 to 7 L/ha) will provide longer term control. The low rate of Roundup should be applied in 50 to 100 L/ha water. If higher water volumes are used add a suitable surfactant. Wait 72 hours before plowing under. Best control of quackgrass is obtained when these herbicides are applied in the fall.
			Roundup WeatherMAX	1.67-8.0 L/ha	12 hours	7	
			Touchdown 480	2.5-7.0 L/ha	12 hours	7	
	14	carfentrazone-ethyl	Aim EC	36.5-117 mL/ha	12 hours	1	Apply in minimum spray volume of 100 L/ha. Refer to label for target weeds, buffer zones and rates. Use high flow rate nozzles to apply the highest spray volume.
Pre-emergence <i>Stale Seedbed Technique</i>	22	paraquat	Gramoxone 200 Sn	2.75-5.5 L/ha	12 hours	-	Apply in 300 – 1100 L of water/ha to foliage of emerged weeds but before the crop has emerged.
		diquat	Reglone 240, Dessicash	2.3-4.6 L/ha	24 hours	-	
Pre-emergence <i>Annual Grasses and some broadleaves</i>	15	propyzamide	Kerb 50 WSP	2.2 kg/ha	24 hours	55	Use on mineral soils only. Do not replant a second crop of lettuce in the same season. One application/year. For transplanted lettuce apply Kerb 7-10 days post-transplant.

Postemergence <i>Inter-row shielded</i>	22	paraquat	Gramoxone 200 Sn	2.75-5.5 L/ha	12 hours	-	Do not spray solution on the crop plant since it could be injured or killed. Use gramoxone for best control of grasses.
		diquat	Reglone 240, Dessicash	2.3-4.6 L/ha	24 hours	-	
	14	carfentrazone-ethyl	Aim EC	36.5-117 mL/ha	12 hours	1	Apply in minimum spray volume of 100 L/ha. Refer to label for target weeds, buffer zones and rates. Use high flow rate nozzles to apply the highest spray volume.
Postemergence	15	propyzamide	Kerb 50 WSP	2.2 kg/ha	24 hours	55	Transplanted lettuce only. Apply 7-10 days after planting. One application per year.
	1	sethoxydim	Poast Ultra plus Merge	0.32-1.1 L/ha 0.5-2 L/ha	12 hours	30	Transplanted lettuce – 1 application per year, between 1- 6 leaf stage Direct seeded – 2 applications/year. First between 1- 6 leaf stage, 2 nd 14 to 21 days after thinning. DO Not exceed 1.1 L/ha/year.
	1	fluazifop-p-butyl and S-isomer	Venture L	2.0 L/ha	12 hours	14	Head & Leaf Lettuce. Apply as a post-emergent broadcast application. One application per season.

Pest	Group	Active Ingredient	Pesticide Product Name	Rate	Restricted-Entry Interval (REI)	Pre-harvest Interval (days)	Remarks
INSECTS:							
Aster Leafhopper	3	cypermethrin	Mako	125 ml/ha	-	14	7-14 day spray schedule. Up to 4 applications/season.
			UP-Cyde	200 ml/ha	12 hours	14	
	1A	carbaryl	Sevin XLR Plus	1.25-2.5 L/ha	12 hours	5 (head lettuce) 21 (leaf lettuce)	Use low rate on young plants. Repeat applications at 7-10 day intervals as necessary. Do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days
	1B	malathion	Malathion 500 EC	1.5-2.75 L/ha	12 hours	3 (head lettuce) 14 (leaf lettuce)	Weekly or twice weekly during growing season.
	-	kaolin	Surround WP	12.5-25 kg/ha	4 hours	-	An OMRI Listed product suitable for organic production. Apply in 500 L of water at 7-14 day intervals once initial infestation is detected. Use high rate for early applications. Do not exceed 25 kg/ha per application.
Aphids	4	imidacloprid	Admire 240 F	10.2 ml/1000plants	12 hours	21	Transplant tray plug drench. Do not follow with soil or foliar treatments
			Admire 240 F	650 ml/ha	24 hours	21	Soil application (field drench) Apply after transplanting in 2000 L water/ha. Max 1 application per year.
			Admire 240 F	200 ml/ha	24 hours	7	Foliar – Do not make foliar application after either a transplant or soil drench of Admire.

	4	acetamiprid	Assail 70 WP Aceta 70 WP	56-86 g/ha	12 hours	7	Max 5 applications per year, alternate with other products.
	1B	acephate	Orthene 75 SP	563-825 g/ha	24 hours	7	Head lettuce only. Maximum of 4 applications/yr
	1B	dimethoate	Lagon 480 EC	700 ml/ha	3 day	7	Leaf lettuce only
	9B	Pymetrozine	FulFill 50WG	193 g/ha	12 hours	14	Apply when aphids first appear. Do not exceed 2 applications/ season. Allow 7 days between applications. Causes feeding cessation shortly after exposure but aphids may remain on the plant for 2-4 days before they die. The use of a non-ionic surfactant at a rate of 0.25% v/v is recommended to improve performance under drought stress conditions.
	29	Flonicamid	Beleaf 50SG	0.12-0.16 kg/ha	12 hours	0 days	Thorough spray coverage of plant foliage is essential. Minimum of 94 L of water/ha. Max 3 applications /season; allow 7 days between applications. Avoid overnight storage of spray mixtures, do not use liquid fertilizer as a carrier; do not use adjuvants.
	4	thiamethoxam	Actara 25 WP	105 g/ha	12 hours	7	Apply before aphids reach damaging levels. Maximum 2 applications/year. Actara is highly toxic to bees, do not apply to blooming crops and wait at least 5 days before placing beehives in a treated field. Allow at least 7 days between applications.
	4C	Sulfoxaflor	Closer SC	100-150 ml/ha	12 hours	3 days	Maximum 2 applications/growing season. Do not apply during crop flowering period or when flowering

							weeds are present in treatment area. Minimum treatment interval = 7 days.
	28	Cyantraniliprole	Exirel	500-1500 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control. For optimum control apply Hasten NT Spray Adjuvant* at an application rate of 0.25% v/v or MSO Concentrate with Leci-Tech* at an application rate of 0.5% v/v. Do not make more than 4 applications per season.
	9D	Afidopyropen <i>NEW 2019</i>	Versys	0.1 L/ha	12 hours	0	<u>Green peach aphid (<i>Myzus persicae</i>), potato aphid (<i>Macrosiphum euphoribae</i>) & lettuce aphid (<i>Nasonovia ribis-nigri</i>).</u> Minimum 7 day interval between applications. Monitor pest population and reapply if necessary once thresholds are reached. Max 4 applications per year.
Aphids and Whiteflies	23	spirotetramat	Movento 240 SC	220-365 ml/ha	12 hours	3	Minimum 7 day interval between applications. Maximum of 730 ml/ha of product applied/season. Movento is TOXIC to bees through direct contamination of pollen and nectar. DO NOT apply during crop flowering period or when flowering weeds are present in the field.
			Movento 150 OD	347-585 ml/ha	12 hours	3	Minimum 7 day interval between applications. Maximum 1.17 L/ha of product applied/season. Movento is TOXIC to bees through direct contamination of pollen and nectar. DO NOT apply during crop flowering period or when flowering weeds are present in the field.

	9D	Afidopyropen <i>NEW 2019</i>	Versys	0.35-0.5 L/ha	12 hours	0	<u>Sweet potato whitefly (<i>Bemisia tabaci</i>) & silverleaf whitefly (<i>Bemisia argentifolii</i>)</u> . Allow a minimum of 7 days between applications. Monitor pest population and reapply if necessary once thresholds are reached. Max 4 applications per year.
Cabbage Looper	11	<i>Bacillus thuringiensis</i>	Dipel 2X DF	275-550 g/ha	-	0	Apply when larvae are small and in the early instar stages. Not effective as larvae become larger.
			Thuricide HPC	2.0-4.25 L/ha	-	0	
	5	spinosad	Success 480 SC	182 ml/ha	12 hours	1	Max 3 applications/yr. Allow 7-10 days between applications
			Entrust 80 W	109 g/ha	12 hours	1	
			Entrust SC	364 ml/ha			
	5	spinetoram	Delegate WG	140-200 g/ha	4 hours	1	Time application with peak egg hatch or small larvae. Repeat applications based on population monitoring. Use higher rate for higher infestations or advanced growth stages. Max. 3 applications/year with a minimum of 5 days between treatments.
	3	cyhalothrin-lambda	Matador 120 EC	83 ml/ha	24 hours	3	Head lettuce only. Maximum 3 applications/year.
	1A	methomyl	Lannate Toss-N-Go	0.51-1.0 kg/ha	24 hours	7	
	1B	acephate	Orthene 75 SP	563-825 g/ha	24 hours	7	Head lettuce only. Max 4 applications per season. Apply in 225-1650 L of water/ha.
28	chlorantraniliprole	Coragen	250 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached. Maximum 4 applications/season. Do not apply more than once every 3 days. Do not exceed 1 L/ha /season. Apply in a finished spray volume of 100L/ha.	

	18	Methoxyfenoziide	Intrepid 240F	0.3 – 0.6 L/ha	12 hours	1	Apply at the first sign of feeding damage or when infestations reach threshold levels as determined by monitoring. Repeat applications after 7-14 days if required based on monitoring. Use higher rate for heavy infestations, advanced growth stages of the target pest or larger crop canopies.
	28	Cyantraniliprole	Exirel	250-500 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control. Do not make more than 4 applications per season.
	11	<i>Bacillus thuringiensis</i> , subsp. <i>aizawai</i> , (Strain ABTS-1857 fermentation solids, spores, and insecticidal toxins)	XenTari WG	500-1000 g/ha	-	0	Apply sufficient spray volume to ensure uniform deposition on all plant surfaces; recommend 500 L per hectare.
Cutworm	3	cyhalothrin-lambda	Matador 120 EC	83 ml/ha	24 hours	3	Head lettuce only. Max 3 applications/year
	3	cypermethrin	Ripcord 400 EC	175 ml/ha	-	21	Do not disturb soil for 5 days after treating
	3	permethrin	Pounce	180-390 ml/ha	12 hours	-	Apply up to the 5-leaf stage. Do not disturb soil for 5 days after treating.
	28	Cyantraniliprole	Exirel	500-750 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control. For early season cutworm control, apply to foliage when rain is not expected in the next 24 hours. For optimal control, apply to smaller plants or when lower portions of plant can receive adequate coverage. Maximum 4 applications/yr.

Tarnished Plant Bug	3	Lambda cyhalothrin	Matador 120 EC	83 ml/ha	24 hours	3	For Head and Leaf Lettuce. Use 500 L/ha of spray volume for thorough coverage. Maximum 3 applications per year, with a 7 day interval between treatments.
	1A	carbaryl	Sevin XLR Plus	2.5-5.25 L/ha	12 hours	5 (head lettuce) 21 (leaf lettuce)	
	4	thiamethoxam	Actara 25 WP	210 g/ha	12 hours	7	Apply before aphids reach damaging levels. Maximum one application/year. This product is highly toxic to bees, do not apply to blooming crops and wait at least 5 days before placing beehives in a treated field.
Brown Marmorated Stink Bug	1B	Malathion	Malathion 85E	1345 mL/ha	-	3 (head lettuce) 14 (leaf lettuce)	Ensure sufficient water volume is used to guarantee thorough coverage. Use a minimum of 500 L water/ha. Apply prior to harvest when treatment thresholds have been reached, as determined by local monitoring.
Onion thrips	5	Spinetoram	Delegate WG	200-336 g product/ha	12 hours	1	Max 3 applications/year. Apply when onion thrips first appear targeting egg hatch and small nymphs. Repeat in 7-10 days if needed.
	23	Spirotetramat	Movento 240 SC	365 ml/ha	12 hours	3 days	Minimum 7 day interval between applications. Maximum allowed per crop season: 730 mL/ha. Apply when thrips are first identified. Movento should be used during the first half of the season when adult populations are relatively low or building. Reductions in numbers of thrips larvae may take 3 to 4 days after Movento is applied.
Armyworm, Beet	28	Cyantraniliprole	Exirel	500 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached.

Armyworm, Fall Armyworm							Thorough coverage is important to obtain optimum control. Make no more than 4 applications per season.
	11	<i>Bacillus thuringiensis</i> , subsp. <i>aizawai</i> , (Strain ABTS-1857 fermentation solids, spores, and insecticidal toxins)	XenTari WG	500-1000 g/ha	-	0	Beet Armyworm. Apply sufficient spray volume to ensure uniform deposition on all plant surfaces; recommend 500 L per ha.
Corn earworm	28	Cyantraniliprole	Exirel	750 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control. Do not make more than 4 applications per season.
Dipteran leafminers (larvae)	28	Cyantraniliprole	Exirel	1000-1500 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control. Do not make more than 4 applications per season.
	4	Acetamiprid	Assail 70 WP Aceta 70 WP	86 g/ha	12 hours	7	Pea leafminer (<i>Liriomyza huidobrensis</i> Blanchard). Begin applications when insect populations reach recognized economic threshold. Thorough coverage is important to obtain optimum control. Do not make more than 5 applications per season. Do not apply more than once every 7 days.
Flea Beetles	28	Cyantraniliprole	Exirel	500-1000 ml/ha	12 hours	1	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control. Re-application interval: 5 days. Max 4 applications/season.

Pest	Group	Active Ingredient	Pesticide Product Name	Rate	Restricted-Entry Interval (REI)	Pre-harvest Interval (days)	Remarks
DISEASES:							
Aster Yellows		-	-	-	-	-	Control Aster leafhoppers
Damping Off	4	metalaxyl	Ridomil 2G	25 kg/ha	-	-	Apply with seed in furrow. Do not use on transplants
	M	thiram	Thiram 75 WP	90 g/25kg seed	24 hours	-	
	4	metalaxyl-M and s-isomer	Apron XL LS	20–40 ml / 100kg of seed	-	-	One application as a seed treatment. Do not apply to leafy greens that are destined to be grown in the greenhouse.
Botrytis (Grey mold)	2	iprodione	Rovral 50 WP	1.5 kg/ha	12 hours	4	Max 4 applications/yr. Start after thinning, apply at 7 day intervals.
	7	boscalid	Cantus WDG	0.285 kg/ha	12 hours	14	Maximum 2 applications/year
	M	ferbam	Ferbam 76 WDG	2 kg/ha	-	-	Spray seedlings before transplant. Apply in 1000L / ha
	-	<i>Bacillus subtilis</i>	Serenade Opti	0.6-1.7 Kg/ha	-	0	Biopesticides that will only suppress the indicated diseases. Begin applications soon after emergence or transplant and continue as necessary on a 7-10 day interval. When conditions are conducive to rapid disease development, use in a rotational program with other registered fungicides. Thorough coverage is essential.
	-	<i>Trichoderma harzianum</i> Rifai strain KRL-AG2	RootShield HC	10 g / L of water	0 hours	-	This biopesticide is for suppression of botrytis blight (<i>B. cinerea</i>). Apply 10 grams/L of water to foliage every 7-14

							days depending upon disease pressure. Apply sufficient spray volume to adequately cover the leaves. This product does not work well when soils are below 10°C.
	7	penthioopyrad	Fontelis	1.25-1.75 L/ha	12 hours	3	Begin applications prior to disease development, continue on a 7-10 day interval. Use higher rate and shorter interval when disease pressure is high. Max seasonal rate is 5.25 L/ha. Do not make more than 2 sequential applications before switching to another mode of action.
Downy Mildew	4M	Metalaxyl-M/mancozeb	Ridomil Gold MZ 68 WP	2.5 kg/ha	24 hours	14	For head lettuce only. Apply every 14 days beginning at thinning. Maximum 3 applications per year.
	U	fosetyl-al	Aliette WDG	2.8 kg/ha	24 hours	7	Max 5 applications/year.
	-	<i>Bacillus subtilis</i>	Serenade Opti	1.7-3.3 Kg/ha	-	0	Biopesticide that will only suppress the indicated diseases. Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.
	44	<i>Bacillus amyloliquifaciens</i> strain D747	Double Nickel LC (1x10¹⁰spores/mL)	5-12.5 L/ha	-	0	Begin applications preventatively when conditions are favorable for onset of disease. Repeat application every 7 - 10 days while conditions favor disease development.
			Double Nickel 55 (5x10¹⁰spores/g)	1-2.5 kg/ha			
40	mandipropamid	Revus plus Non-ionic adjuvant	400-600 ml/ha plus 0.125% v/v	12 hours	1	For control of Downy Mildew and Blue Mold. Applications should begin prior to disease development and continue throughout the season on a 7-10 day schedule of fungicides, following resistance management guidelines. Maximum 4 applications per year.	

	43	fluopicolide	Presidio	220-292 ml/ha	12 hours	2	Apply in 200-1000 L/ha. For resistance management, Presidio must be tank-mixed with a labelled rate of another fungicide registered for the target pathogen with a different mode of action. Apply in a tank mix with Aliette. Follow the most restrictive use directions of either label. Make foliar applications on a 7-10 day schedule beginning with initial flowering, or when disease conditions are favourable, but prior to disease development. Use lower rate and longer interval as preventive applications. Use the higher rate and shorter interval if disease is present.
	40	Dimethomorph	Acrobat 50 WP	450 g/ha	2 days (hand harvesting) 12 hours (no entry)	0	Must be applied as a tank mix with another fungicide active against downy mildew. Begin applications when conditions favour disease development. Continue applications on a 5-10 day schedule as needed until weather conditions favouring infection and sporulation decrease. DO NOT use less than 200 L of water/ha for ground applications. Max 5 applications/year.
	33	Mono and dibasic sodium and potassium and ammonium phosphites	Phostrol	2.9-5.8 L/ha	12 hours	1	For preventative control of downy mildew , begin foliar applications when conditions favouring disease development exist and continue on a 7-21 day interval, if needed. Max 7 applications per year.
	40,45	Ametoctradin, dimethomorph	Zampro	1.0 L/ha	Hand harvesting – 1 day All other	0	To reduce the risk of development of fungicide resistance use in rotation with other fungicides having a different mode of action active against downy

					activities – 12 hours		mildew. Begin applications prior to disease development and continue on a 5-7-day interval. Max 3 applications.
	21	Cyazofamid	Torrent 400 SC + Sylgard 309	0.20 L per hectare	12 hours	0	Make the first application after seeding or once the lettuce transplants are set in the field. Subsequent applications should be initiated when weather conditions are favourable for downy mildew development or when disease first appears in the field. All subsequent applications on a 7-14 day schedule, using the shorter interval when disease pressure is moderate to severe. Max 6 applications/ season. A plant back interval of 30 days is required.
	U15	Oxathiapiprolin	Zorvec Enicade	0.175- 0.35 L/ha	12 hours	0	Begin applications prior to disease development and continue on a 5-14 day interval. Use higher rate and shorter interval when disease pressure is high.
	19	Polyoxin D zinc salt <i>NEW 2019</i>	Diplomat 5SC	463 – 926 ml/ha	-	0	Apply as a foliar spray in sufficient water to provide thorough coverage of foliage. Begin as a preventative application when conditions favour disease development and continue on a 7-14 day interval as needed to maintain suppression. Do not apply more than 150 g a.i./ha/season or 12 months.
Sclerotinia (Head and Leaf Drop)	7	boscalid	Cantus WDG	0.385 kg/ha	12 hours	14	Max 2 applications/year.
	44	<i>Bacillus subtilis</i>	Serenade Opti	0.6-1.7 Kg/ha	-	0	Biopesticide that will only suppress the indicated diseases. Make the first application at planting (or immediately following planting but prior to crop emergence). Make a 2 nd application as a directed spray with multiple nozzles per seed line in sufficient water to

							ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Repeat applications on 10-14 day intervals if conditions for disease development persist.
	44	<i>Bacillus amyloliquefaciens</i> strain D747	Double Nickel LC (1x10¹⁰spores/mL) Double Nickel 55 (5x10¹⁰spores/g)	Foliar: 5-12.5 L/ha Low disease pressure: 1-5 L/ha Soil: 1-2.5 L/ha Foliar: 1-2.5 kg/ha Low disease pressure: 0.2-1 kg/ha Soil: 0.2-0.5 kg/ha	-	0	Growth stage: From planting to formation of the head of lettuce. Repeat application every 3 to 10 days for as long as conditions favor disease development. See Soil application on label. Apply at or after planting (but before plant emergence) as a banded seed line treatment, see Banded/in furrow application on label. Alternatively, apply to seed bed via overhead sprinkler chemigation with germination water, see Chemigation Instructions on label. Make second application at thinning or cultivation in sufficient water and multiple nozzles to ensure thorough coverage of lower leaves and surrounding soil surface. Incorporation with light irrigation after application may improve disease control. Repeat at 10-14 day intervals if conditions promoting disease persist.
	7	penthiopyrad	Fontelis	1.25-1.75 L/ha	12 hours	3	Begin applications prior to disease development; continue on a 7-10 day interval. Use higher rate and shorter interval when disease pressure is high. Max seasonal rate is 5.25 L/ha. Do not make more than 2 sequential applications before switching to another mode of action.

	7	Isofetamid	Kenja 400SC	0.90 L/ha	12 hours	14	On direct seeded lettuce make the first application after emergence, thinning or prior to onset of disease development On transplanted lettuce make the first application immediately after transplanting or prior to the onset of disease development. Make a second application 14 days later if conditions continue to favor disease development. Do not make more than 2 applications per year.
Powdery Mildew	-	<i>Bacillus subtilis</i>	Serenade Opti	1.7-3.3 Kg/ha	-	0	Biopesticides that will only suppress the indicated diseases. Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.
	13	quinoxifen	Quintec	240 ml/ha	12 hours	1	Maximum of 4 applications per year. Repeat applications on 10-14 day intervals. Apply before visual symptoms of Powdery Mildew appear. Quintec has no curative properties and will not control established infections. * Note * – product is toxic to bees exposed to direct treatment, drift, or residues on crops or weeds.

Use the following web link to search for any pesticide label mentioned in this guide, or any other pesticide registered in Canada:

<http://pr-rp.hc-sc.gc.ca/lr-re/index-eng.php>

PESTICIDE EMERGENCY CONTACT INFORMATION

Poison Control Centres		
Nova Scotia	800.565.8161 or 902.428.8161	IWK, Halifax, NS
New Brunswick	911	Ask for Poison Information
Prince Edward Island	800.565.8161 or 902.428.8161	IWK, Halifax, NS
Newfoundland	709.722.1110	Dr. Charles A. Janeway Child Health Care Centre, St. John's, NF

Environmental Emergencies (Pesticide Spills)	
Transport Canada Regional Operations Centre (24 hours)	
Nova Scotia	800.565.1633
New Brunswick	800.565.1633
Prince Edward Island	800.565.1633
Newfoundland	800.563.9089

ABBREVIATIONS & CONVERSIONS

Formulation and Measurement Abbreviations			
FORMULATIONS		MEASUREMENTS	
DF	Dry flowable	mL	millilitre
EC, E	Oil-based emulsifiable concentrate	kPa	kilopascal
EW	Water-based concentrate	kg	kilogram
EG	Water dispersible granule	g	gram
F	Flowable	L	litre
G	Granule	BIU	Billions of International Units
SC	Suspension concentrate	ppm	parts per million (1000 ppb)
Sn	Solution	ppb	parts per billion (1/1000 ppm)
SP	Soluble powder		
WDG	Wettable dry granule		
WP, W	Wettable powder		

Helpful Conversions¹	
kPa X 0.14 = pounds per square inch (psi)	millilitres X 0.035 = fluid ounces
hectares X 2.47 = acres	litres X 35 = fluid ounces
kilograms X 2.2 = pounds	litres X 0.22 = imperial gallons
kilograms per hectare X 0.89 = pounds per acre	litres per hectare X 14.17 = fluid ounces per acre
kilograms per hectare X 0.40 = kilograms per acre	litres per hectare X 0.40 = litres per acre
	degree-days C X 1.8 = degree-days F

¹ **Pesticide Units of Measurement**

It is not recommended to convert label rates to imperial units because there is a high probability of mathematical and rounding errors. Present day pesticides are formulated to be more effective in smaller amounts. Therefore, even small conversion errors can lead to the use of incorrect rates (either too high or too low). Use metric – you will be glad you did!