

## Order of Operations at June Bearing Strawberry Renovation:

Decide what stays and what needs to be quickly terminated

- if terminating apply Roundup plus broad spectrum insecticide

Apply herbicide

Lontrel or 2,4-D

2,4-D .6l/ac ( 2-3 days Mow)

Lontrel XC 120 ml/ac (7-10 days Mow)

5 days later Mow

Fertilizer

50-60 units of nitrogen + P + K

Refrain from using 0-0-60 as a potassium source

Apply AgriMek for cyclamen mite?

Agri Mek 95 ml/ac in 300 l/ac

(Maximum 2 apps/year)

surfactant at 0.2% ag surf

apply about 2 weeks after previous herbicide application

5 days after Agrimek application Sinbar in rain

280-340 grams/ac

10 days after Sinbar apply grass herbicide

(Poast, Select, Venture)

Aphids?

- allow for 10-14 days between Sinbar and 2,4-D or Lontrel application

-allow 10-14 days between Sinbar and Venture

- allow 5-7 days between Sinbar and Agri mek application

Control powdery mildew, common leaf spot

Captan/Meastro for Anthracnose

Other management steps that some growers use:

Apply Aliette before herbicide to restore root health

Apogee for runner control if too many runners?

Chain harrow to remove older crowns and thin within row

Narrow the rows with ignite? Controls weeds also between rows?

ripping using deep tillage to improve drainage an alleviate compaction